

DOLUTEGRAVIR INTERACTIONS INFORMATION SHEET

POTENTIAL INTERACTIONS WITH DOLUTEGRAVIR (DTG)

Interacting drug	Effect of co-administration	Recommendation
Rifampicin	↓ DTG	Double DTG dose to 50 mg 12-hourly. If on TLD FDC, add DTG 50 mg 12 hours after TLD dose
Anticonvulsants: Carbamazepine Phenobarbital Phenytoin	↓ DTG	Avoid coadministration if possible. Alternative agents that do not interact with DTG include valproate, lamotrigine, levetiracetam, and topiramate. Remember that valproate is contra-indicated during pregnancy. Double DTG dose to 50 mg 12-hourly for carbamazepine if an alternative anticonvulsant cannot be used
Metformin	↑ Metformin	DTG increases metformin levels. Maximum metformin dose 500 mg 12-hourly
Polyvalent cations (Mg²⁺, Fe²⁺, Ca²⁺, Al³⁺, Zn²⁺) e.g. antacids, sucralfate, multivitamin and nutritional supplements		
Calcium	↓ DTG	Interaction occurs if taken together on empty stomach. To prevent this, take at the same time, with food. Without food, take the calcium supplement a minimum of 2 hours after or 6 hours before DTG
Iron	↓ DTG	Interaction occurs if taken together on empty stomach. To prevent this, take at the same time, with food. Without food, take the iron supplement a minimum of 2 hours after or 6 hours before DTG Remember: Take calcium and iron at least 4 hours apart
Magnesium/ aluminium containing antacids	↓ DTG	We do not know if food intake eliminates this interaction. Antacids should be taken a minimum of 2 hours after or 6 hours before DTG
Sucralfate/Zinc	↓ DTG	We do not know if food intake eliminates this interaction. Sucralfate/zinc should be taken a minimum of 2 hours after or 6 hours before DTG



MEDICINES
INFORMATION
CENTRE

This publication was supported under funding provided by the Global Fund to Fight AIDS, Tuberculosis and Malaria through the National Department of Health of South Africa and the NDoH Pharmacovigilance Centre for Public Health Programmes. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the Global Fund or the National Department of Health of South Africa



NEED HELP?

Contact the TOLL-FREE National HIV & TB Health Care Worker Hotline

0800 212 506 / 021 406 6782

Alternatively "WhatsApp" or send an SMS or "Please Call Me" to 071 840 1572
www.mic.uct.ac.za

Download our free App: SA HIV/TB Hotline



Published October 2020, Version 1

IF YOU TAKE DOLUTEGRAVIR (e.g. TLD) AND NEED TO TAKE CALCIUM AND IRON SUPPLEMENTATION (e.g. DURING PREGNANCY) DO IT LIKE THIS:

WITH FOOD



08:00 TLD + Calcium + FOOD



13:00 Iron supplement



20:00 Calcium

Antacid can be taken after 10h00 and before 02h00, but not at the same time as iron. Try to separate the dosing of iron and antacids as much as possible

WITHOUT FOOD



06:00 TLD
08:00 Calcium



13:00 Iron supplement



20:00 Calcium

Antacid can be taken after 8h00 and before 24h00, but not at the same time as iron. Try to separate dosing of iron and antacids as much as possible